

2023 Recommended Immunizations for Children from Birth Through 23 Months

VACCINE	Birth	1 MONTH	2 MONTHS	4 MONTHS	6 MONTHS	12 MONTHS	15 MONTHS	18 MONTHS	19-23 MONTHS
HepB Hepatitis B	HepB	HepB			HepB				
RV* Rotavirus			RV	RV	RV*				
DTaP Diphtheria, Pertussis, & Tetanus			DTaP	DTaP	DTaP		DTaP		
Hib* <i>Haemophilus influenzae</i> type b			Hib	Hib	Hib*	Hib			
PCV13, PCV15 Pneumococcal disease			PCV	PCV	PCV	PCV			
IPV Polio			IPV	IPV	IPV				
COVID-19** Coronavirus disease 2019					COVID-19**				
Flu† Influenza					Flu (One or Two Doses Yearly) †				
MMR Measles, Mumps, & Rubella						MMR			
Varicella Chickenpox						Varicella			
HepA‡ Hepatitis A						HepA‡		HepA‡	

FOOTNOTES

RV* **Hib***
Administering a third dose at age 6 months depends on the brand of Hib or rotavirus vaccine used for previous dose.

COVID-19** Number of doses recommended depends on your child's age and type of COVID-19 vaccine used.

Flu† Two doses given at least 4 weeks apart are recommended for children age 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group.

HepA‡ Two doses of Hep A vaccine are needed for lasting protection. The 2 doses should be given between age 12 and 23 months. Both doses should be separated by at least 6 months. Children 2 years and older who have not received 2 doses of Hep A should complete the series.

ADDITIONAL INFORMATION

1. If your child misses a shot recommended for their age, talk to your child's doctor as soon as possible to see when the missed shot can be given.
2. If your child has any medical conditions that put them at risk for infection (e.g., sickle cell, HIV infection, cochlear implants) or is traveling outside the United States, talk to your child's doctor about additional vaccines that they may need.

Talk with your child's doctor if you have questions about any shot recommended for your child.